



Cite this article: Noble LM, Andrianopoulos A. 2013 Reproductive competence: a recurrent logic module in eukaryotic development. *Proc R Soc B* 280: 20130819. <http://dx.doi.org/10.1098/rspb.2013.0819>

Received: 4 April 2013

Accepted: 14 June 2013

Subject Areas:

developmental biology, genetics, evolution

Keywords:

development, reproduction, competence, vegetative phase change, life history, fungi

Author for correspondence:

Luke M. Noble

e-mail: luke.noble@gmail.com

†Present address: Center for Genomics and Systems Biology, Department of Biology, New York University, New York, NY 10012, USA.

Reproductive competence: a recurrent logic module in eukaryotic development

Luke M. Noble[†] and Alex Andrianopoulos

Department of Genetics, University of Melbourne, Victoria 3010, Australia

Developmental competence is the ability to differentiate in response to an appropriate stimulus, as first elaborated by Waddington in relation to organs and tissues. Competence thresholds operate at all levels of biological systems from the molecular (e.g. the cell cycle) to the ontological (e.g. metamorphosis and reproduction). Reproductive competence, an organismal process, is well studied in mammals (sexual maturity) and plants (vegetative phase change), though far less than later stages of terminal differentiation. The phenomenon has also been documented in multiple species of multicellular fungi, mostly in early, disparate literature, providing a clear example of physiological differentiation in the absence of morphological change. This review brings together data on reproductive competence in Ascomycete fungi, particularly the model filamentous fungus *Aspergillus nidulans*, contrasting mechanisms within Unikonts and plants. We posit reproductive competence is an elementary logic module necessary for coordinated development of multicellular organisms or functional units. This includes unitary multicellular life as well as colonial species both unicellular and multicellular (e.g. social insects such as ants). We discuss adaptive hypotheses for developmental and reproductive competence systems and suggest experimental work to address the evolutionary origins, generality and genetic basis of competence in the fungal kingdom.

1. Introduction

Differentiation is the acquisition of specialized capabilities, and development is the overarching ontogenesis. Development unfolds through interplay between the genome, constrained by its evolutionary history, and the environment, to which the genome is adapted. Generation of distinct cell types, a conveniently discrete criterion, is typically considered the minimal unit. Development is induced, meaning simply it is caused by something, and the causes are said to be inducers. Organisms are exposed to the environment over very different scales, however, from the extreme intimacy of a single-celled bacterium to a highly differentiated, homeostatic, long-lived animal. It follows that in more exposed organisms, development is directly induced by certain environmental cues, whereas in more insulated organisms, induction is effected by endogenous cues (hormones) during ontogenesis. Part way along this continuum, many relatively simple organisms use both secreted signals and environmental induction to coordinate development. The response to a fluctuating environment may vary according to homeostatic capacity but is necessarily limited by the metabolic economy (energy), the ultimate determinant of proliferative and reproductive capacity.

In addition to induction and the subsequent response, another fundamental but neglected property of development is competence: the ability to respond to induction [1–3]. In this sense, the term ‘competence’ was introduced into the English scientific literature in 1932 by Waddington, in studying how the vertebrate notochord induces ectoderm to become neural rather than epidermal tissue [4,5]. Waddington outlined three separable aspects of development: competence, the ability to respond to induction; evocation, the inducing stimulus itself and individuation, the resulting process of differentiation.

As for the general terms of differentiation and development, competence can be recognized at varying levels of complexity, from the subcellular




Figure 1. *Aspergillus nidulans* reproductive cycles and the logic of competence. (a) Depiction of the four reproductive cycles [13]—asexual, sexual, parasexual (producing recombinant haploid nuclei) and hyphal growth (producing clonal nuclei, although replication errors may contribute to heterokaryosis in a mycelium of sufficient size)—all of which may occur simultaneously in a single colony. (b) Developmental timing in *A. nidulans* for a single, isolated conidium growing at 37°C in the presence of excess, preferred nutrients [12]. (c) Defining competence in *A. nidulans* [14]. Conidiophores per colony plotted over time since inoculation, in response to variable time of induction (black arrows). Development was induced by exposure of single spore-derived colonies to air and light. Minimum time to conidiation (blue arrow) comprises competence acquisition (grey arrow) and a constant maturation period for conidiophore production (red arrows). Vegetative growth continues indefinitely in the absence of induction and the presence of sufficient nutrients.

(e.g. cell cycle [6] or gene regulatory network [7,8]) to the organismal (e.g. reproductive competence) [9–12]. ‘Competence’ has therefore been applied in different contexts to describe many different processes. Developmental competence could be considered as any measurable state that must precede another in the development of an organism, organ or cell, but a definition encompassing all of ontogenesis at all scales is not useful. More so is the characterization of developmental competence as a ‘checkpoint’ between discrete life-history phases, such as metamorphosis and reproductive maturity.

2. Fungal competence

Acquisition of reproductive competence in the model filamentous Ascomycete *Aspergillus nidulans* is a necessary prelude to asexual development, producing dormant spores, and is associated with rapid change in diverse biochemical systems. The existence of competence has also been established in the related pathogen *Aspergillus fumigatus* and in various *Penicillium* and *Trichoderma* species, suggesting conservation in filamentous Ascomycetes at least.

(a) Reproductive competence in *Aspergillus nidulans*

Aspergillus nidulans reproduction involves some combination of vegetative growth (clonal nuclei distributed throughout the

mycelium), asexual development (uninucleate conidiospores), parasexual development (recombinant nuclei via a diploid stage) and sexual development (ascospores via meiosis; figure 1a). Conidia break dormancy in the presence of water and an assimilable carbon source with a period of isotropic growth and nuclear division, followed by establishment of a polar growth focus and emergence of a hyphal germ tube [15–18]. Nuclear division is synchronous over the first three to four mitoses (influenced by culture conditions and growth rate), and the first incomplete hyphal barrier, or septum, is laid down around this time [19,20]. Nuclear division within the mycelium subsequently becomes asynchronous, but mitosis and biomass accumulation remain exponential given sufficient nutrients. Actively growing apical septate compartments are highly enriched in nuclei but are also much larger than intercalary compartments, so that the ratio nuclei to cytoplasm remains similar—known as the hyphal growth unit (G). Subsequent subapical branching and mutual hyphal avoidance (negative autotropism) due to accumulation of secreted morphogens leads to the characteristic fungal network, with exploratory growth at the colony exterior and a developmentally committed interior mycelium [21].

Developmental genetic networks have been studied extensively, especially those underlying asexual development, although integration of conserved signal transduction pathways such as *ras/cAMP/PKA* and G-proteins with fungal-specific

transcription factors is rudimentary [22–24]. Diverse inputs ultimately converge on a terminal network comprising transcription factors such as bristle (BrlA) and abacus (AbaA), which establish commitment through feedback loops in the developing conidiophore and drive production of new cell types required for spore generation [25–27].

Without exposure to an inducing air interface and light, however, development is repressed, given nutrient sufficiency. While external nutrient limitation is not a primary inductive cue for differentiation, self-imposed limitation may play a crucial role. Studies of differential gene expression and enzyme activity during conidiation in Ascomycetes demonstrate distinct metabolism in the colony interior and reproductive structures [28–32]. Developmental gene expression is spatially restricted, and ectopic expression of key regulators results in generalized loss of nutrient acquisition and growth arrest [33,34].

Axelrod [12] recognized a minimum of 20 h growth at single colony density is required before *A. nidulans* becomes competent to respond to induction of sporulation (figure 1b). Once competent, the period required for production of a conidiophore from vegetative mycelia (known as maturation) is a minimum of approximately 4.5 h at 37°C (figure 1c).

In subsequent work, a number of biochemical correlates of competence acquisition were identified. First, the inducibility of two enzymes, nitrate reductase and extracellular protease, falls sharply coincident with developmental competence, while activity of a constitutive enzyme in the pentose phosphate pathway, glucose-6-phosphate dehydrogenase, does not [35]. Thus, it appeared that a change in developmental potential was associated with a pleiotropic change in inducibility of enzymes not themselves required for morphological development. While generalization based on this small sample is fraught with potential for over interpretation, it is possible such a change might occur due to altered transcriptional potential (e.g. chromatin state), protein activity or membrane transport.

Second, intracellular cAMP levels fall sharply during early growth [36]. A role for cAMP-dependent PKA in metabolic activation of spores for diverse fungi is known [16,17,37], and one hypothesis for competence timing is a cAMP cycle determined by conidial deposition. Functions of this conserved chemical messenger in morphogenesis of non-yeast fungi are not well delineated, but gross defects in cell-cycle, sporulation, secondary metabolism, pathogenesis and nutrient sensing have been reported in a range of species [38–40].

Third, media replacement and levels of glucose as carbon and of ammonium or glutamate as nitrogen have no effect on the timing of competence acquisition, although rate of conidiophore production is clearly affected [41]. Competence is therefore stable against external nutrient levels, both high and low, although utilization of non-preferred carbon sources does alter timing (L. M. Noble, A. McLauchlan, K. Hubner, A. Andrianopoulos 2013, unpublished data).

Fourth, and most enigmatic, while acid extraction of competent mycelium results in total hydrolysis of DNA, the same procedure applied to precompetent mycelium results in near complete resistance of DNA to hydrolysis [42]. This result was traced to the interfering action of ferric iron in the mycelial lysate, suggesting ‘an alteration in the concentration, compartmentalization, or species, of iron binding components’ [42]. In all, these gross changes in diverse processes suggest competence acquisition is a time of systemic transition, but do not yet coherently implicate any single underlying cause or mechanism of transduction.

(b) *Aspergillus nidulans* competence mutants

Axelrod and others employed a simple screen to identify *precocious* competence mutants [12,14]. Subsequent studies showed the effect size in two (genetically uncharacterised) mutants was proportional to glucose uptake rate [43,44]. Specifically, precociousness correlated with both maximum rate of glucose uptake after germination and with the rate of transport decline as competence approached. Fructose, sucrose, alanine and the non-metabolizable glucose analogue 2-deoxyglucose showed a similar uptake rate decline in wild-type, whereas acetate did not. The carbon catabolite derepression system, which hierarchically controls utilization preferences, also differed markedly between states. Precompetent mycelium grown on glucose was derepressed for fructose transport by starvation more rapidly and to a maximal level four to five times higher than that of competent mycelium.

In sum, this work illustrated a relationship, not necessarily a causal one, between nutrient uptake and competence. It is nonetheless tempting to speculate that competence in fungi, as in other organisms, involves a nutrient accumulation threshold. Axelrod’s [12] observation that inhibition of protein synthesis produces a dose-dependent delay in competence, yet has little effect (initially) on the subsequent production of conidiophores indeed suggests a greater level of developmental autonomy after competence acquisition.

While no *precocious* genes have been cloned, a second class of competence mutants, temperature sensitive *aconidials*, identified multiple subunits of the COP9 signalosome (CSN), a highly conserved protein complex with homology to the proteasome and eukaryotic initiation factor 3 complexes [45–49]. Mutations affecting ras protein activity also modulate, positively and negatively, competence timing and other developmental transitions [50].

(c) Ascomycetes: *Trichoderma* and *Penicillium* species

Trichoderma (*Hypocrea*) species have a long history of study, primarily due to antagonistic relationships with plant pathogenic fungi. Developmental competence in this genus refers to two, possibly distinct, phenomena. First is the variable ability of hyphae within a colony to respond to photoinduction of conidiation [51]. The second, unstudied beyond its initial description, is directly analogous to that of *Aspergilli* in that a minimum period of growth is required before ‘accepting’ photoinduction (figure 2a) [52].

Research into secondary metabolite production in *Penicillium* species found that sporulation could be induced in liquid culture under certain conditions [56]. The addition of calcium ions was able to fully mimic an inducing air interface, and for *Penicillium notatum*, ‘initial sporulation time varied inversely with... the inoculum load’ (figure 2b; [53, p. 419, 57,58]). ‘Mature’ medium, which had supported growth of various fungi, advanced competence acquisition, in some cases to a greater extent than that from *P. notatum* itself.

This body of research led to isolation from *Penicillium cyclopium* of the only known fungal competence hormone [59]. Related diterpenoid molecules were identified, probably derived from the mevalonate pathway responsible for key hormones across all kingdoms of life, named conidiogenol and conidiogenone (figure 2c). Although calcium lowered the effective threshold, conidiogenone alone was able to accelerate competence acquisition and also, surprisingly, induce sporulation. This suggests both competence and induction are




Figure 2. Reproductive competence in *Trichoderma* and *Penicillium* species. (a) The relationship between age and conidiation capacity in *Trichoderma viride* [52]. At left, colonies grown on filter papers were fixed and stained with cotton blue at time of induction by light. At right, duplicate photoinduced colonies after a further 40 h incubation in the dark. A ring of conidia is visible at the colony fringe for colonies of age 20 h or greater. (b) Identification of density dependence for development of *Penicillium notatum* (*chrysogenum*) [53]. Inoculum load (conidia per ml of medium) is plotted against sporulation time (induced by addition of calcium) and dry weight at time of sporulation. A minimum of 17.5 h is required under these conditions for sporulation. In other experiments, it was found that 6 h is required for conidiophore production (maturation), regardless of inoculum load. (c) The mevalonate pathway produces key primary and secondary metabolites in diverse species [54]. The diterpenoid competence hormone, conidiogenone, from *Penicillium cyclopium* is an effector of both competence acquisition and conidiation induction in this species [55].

hormone thresholds, reporting organism density and emergence into an aerial environment where diffusion is limited [55].

3. Competence by any other name?

Notable work on reproductive competence has been carried out in plants (vegetative phase change; [60–62]) and mammals [63–67]. The amoeba *Dictyostelium discoideum* is a special case, with multiple competence thresholds identified at cellular and multicellular levels throughout its experimentally tractable life history. We next discuss competence systems in plants and *Dictyostelium* occupying an equivalent life-history stage to fungal reproductive competence.

(a) Vegetative phase change in plants

Most developmental research in spermatophyte plants has focused on genetic networks that regulate flower development. Between germination and seed production, however, plants also switch from juvenile to morphologically distinct, reproductively competent, mature vegetative growth [11,60,62]. Phase change is particularly obvious in woody plants, such as English ivy and some cacti, where the juvenile phase is protracted and distinctive in growth form (figure 3a,b) but was perhaps first documented in the course of fruit-tree grafting [68]. It has now been studied at the molecular level in *Arabidopsis*, maize and other species, addressing the measurement of time during development and the environmental and endogenous signals involved in regulating transitions.

Basic factors such as photoperiod and temperature, critical as they are to carbon metabolism and environmental forecasting, play an important role in plant competence acquisition [71]. The diterpenoid gibberellin is another important factor. Although genetic networks peripheral to this hormone in phase change are not well understood, biosynthetic and hypersensitive mutants show altered onset of maturity [72,73]. Most illuminating, numerous studies have implicated a conserved role for RNA silencing [74–77]. MicroRNA miR156 has been identified as a crucial control point through targeting the *SPL* (SQUAMOSA promoter-binding protein-like) family of transcription factor genes [78]. These factors, in turn, enable flowering and establish mature epidermal identity by activating further activators and repressors of repressors, including miR172 [61,77]. The inverse expression pattern of these two miRNAs, with miR156 abundant in juvenile shoots but decreasing over time, and negative feedback from their cognate transcription factor targets, underlies the bistable morphologies and reproductive capabilities associated with the phases (figure 3c). Recent work suggests that sugars regulate miR156 levels [79], and that gibberellin and miR156 signalling may be integrated at the level of *SPL* factor regulation [80], providing a direct link between central carbon metabolism and phase change.

(b) Dissecting *Dictyostelium* development

Historically grouped with fungi, soil-dwelling slime moulds are now placed within a sister clade to the Opisthokonts,




Figure 3. Vegetative phase change and transcriptional regulatory networks. Two examples of distinct morphologies associated with vegetative phases in plants. (a) Growth habit and leaf shape in English ivy (*Hedera helix*) for juvenile (lower) and adult (upper) phases [68]. (b) *Melocactus intortus* juvenile body (lower) and (upper) the adult apical cephalium [69]. (c) Conserved transcriptional networks controlling phase change in plants [70]. Sugars regulate levels of microRNA miR156, which in turn determines expression of the SPL family of transcription factors required for phase change and flowering.

the Amoebozoa [81], and most recent phylogenomic analyses merge these into a monophyletic supergroup, the Unikonts [82]. As for plants and fungi, the earliest stages of ontogenesis for *D. discoideum* most relevant to reproductive competence acquisition have received little attention relative to the later spectacular transition to multicellularity for which the model organism is most famous. *Dictyostelium discoideum* is in fact highly derived among Amoebozoa [83], many of

which develop only by encystment or production of very simple fruiting bodies—several clades produce a spore and crude stalk from a single cell. As for prokaryotes with similar reproductive modes, such as myxobacteria [84–86], development in *Dictyostelium* is a direct response to starvation and high population density, mediated by quorum signals and other secreted morphogens [87–90]. cAMP released from amino acid-starved cells above a critical density settles into

synchronized oscillations throughout the population, inducing polarization and chemotaxis towards a central organizing centre. After sufficient exposure, cells become 'aggregation competent' [91], forming a migratory slug that searches for favourable fruiting conditions and, eventually, an upright structure promoting spore dispersal.

Competence to respond to starvation is dependent on constitutively produced autocrine signals [92]. At least three are known, termed prestarvation factors (PSFs), though the responsible genes are not. At a threshold PSF: bacteria (food) ratio, production of another morphogen, conditioned medium factor, leads to transcriptional induction of a small number of pivotal genes, including those for the cAMP receptor and a heterotrimeric G-protein through which cAMP signalling is propagated [93].

Interestingly, this competence threshold is not always absolute—spores of at least one species are able to directly re-enter development without a period of feeding, owing to an unusual positive relationship between spore density and germination (the relationship is negative in other *Dictyostelids* and in many fungi). This could be reiterated, producing successive generations of progressively smaller cell size, indicating spore metabolic reserves are not strictly limiting [94].

4. Questions of mechanism and generality

A threshold governing reproductive capacity is a common feature of development, whether of prokaryotes or eukaryotes, single-celled or multicellular. In *D. discoideum*, competence checks govern multiple transitions from a population of single cells to a cohesive multicellular entity. Quorum sensing molecules (typically divergent, with the notable exception of cAMP) are channelled into conserved signal transduction pathways (e.g. G-proteins) to coordinate behaviour. This includes the earliest stage of vegetative growth, comparable with competence in fungi, where signals reporting organism density and nutrient availability are integrated to determine the switch from niche exploitation to reproduction and dispersal. Competence thresholds in plants and other Unikonts are mediated by intercellular signals and nutrient acquisition. The apparent broad conservation of mechanisms invites several questions and suggests lines of enquiry to compare basal genetic underpinnings, within fungi and more broadly.

(a) How does fungal reproductive competence relate to known mechanisms of spatio-temporal differentiation?

Metabolism in fungi is often discussed as a singular phenomenon, research being grounded on homogeneous liquid culture and unicellular yeasts. Classical studies of the filamentous growth form, developmental studies (which usually must embrace heterogeneous solid culture) and research aimed directly at measuring spatial differentiation within the mycelium illustrate a different picture [15,19,29,95,96]. This may be of relevance for competence acquisition, with intra-hyphal specialization of anabolism within the mycelium interior, and catabolism, secretion and sensory integration at the hyphal apex [21]. The relationship between this hypothesis and established mechanisms of polarity establishment and maturation during early growth remains to be clarified [20,97–100]. Among early changes in hyphal differentiation

are the demarcation of hyphal compartments by septa and establishment of apical-high gradients of reactive oxygen species, Ca^{2+} and protons [96,97,101,102]. It would be of interest to monitor these landmarks temporally to see if their establishment coincides with competence acquisition or is altered in mutants.

For sessile organisms on solid substrate, autoinducers accumulate as a predictable function of density and distribution and report physical aspects of the environment such as diffusion [103]. Competence in fungi is density dependent, even when nutrients are maintained in excess [41]. This presumably means that multiple spores can more rapidly form a mycelium equivalent to that of an older colony derived from a single spore, and so density approximates age. A large and growing number of autoregulatory substances secreted by mycelial fungi are known, including oxylipins—lipidic signalling molecules found across all eukaryotic kingdoms [104–107]—fungal-specific secondary metabolites [108] and growth by-products such as bicarbonate ([92], reviewed in [105]). With the exception of the diterpenoid conidiogenone produced by *P. cyclopium* [59], none are known to alter competence timing. Competence in fungi may be sensitive to both specific morphogens and general effects of environment modification as an inevitable consequence of growth, such as pH.

(b) How is nutrient acquisition related to competence acquisition?

Evidence for reproductive competence acquisition as a nutrient accumulation threshold across diverse organisms is strong. In humans, fat and leptin levels predict sexual maturity better than chronological age [63,65,109]. Many plants require a critical biomass accumulation before flowering can be induced [110]. Age at maturity is negatively correlated with growth rate in insects [111], and similarly, moulting and metamorphic competence is governed by a critical weight threshold [112–114].

Competence in *A. nidulans* is indeed dependent on metabolism based on (i) correlation with glucose uptake rate, (ii) temperature dependence and (iii) slower acquisition on non-preferred carbon sources (although excess nutrients do not accelerate competence, suggesting metabolic saturation). However, the threshold does not appear to represent some absolute minimal attainment of nutrients for successful reproduction. *Penicillium cyclopium* exposed to the competence hormone conidiogenone from inoculation is able to undergo 'microcycle' conidiation shortly after germination [59], and the dictyostelid *Dictyostelium stoloniferum* is able to cycle through multiple generations without intervening growth [94]. Reproductive competence thresholds even in relatively simple eukaryotes may therefore represent constrained but variably plastic, optimized trade-offs rather than absolute limits.

(c) Is competence acquisition always adaptive?

Maintenance of discrete growth phases in plants is considered adaptive [60,68,115], citing, for example, expression of distinct resistance mechanisms where predators and parasites vary between phases as a plant colonizes the aerial niche [116]. Adaptive hypotheses have also been proposed for metamorphic competence in marine invertebrates [117–119] and holometabolous insects [120], and sexual maturity in

mammals [9]. This may be true, but may also be no more than *post hoc* justification according to ecology, without bearing on why a competence state first arose.

Life forms adapt to their environment, within the genetic, morphological and behavioural constraints set by population dynamics and evolutionary history. We suggest reproductive competence is an elementary logic module, which by necessity has been elaborated upon to allow coordinated development of multicellular organisms or functional units. This includes unitary multicellular life as well as colonial species both unicellular and multicellular (e.g. social insects such as ants, which reproduce colonies only above a critical size [121]). Such behaviour would be expected to emerge from nonlinear dynamics of genetic networks reflecting strong preference for stable, committed development.

The simplest hypothetical ancestral state exhibiting reproductive competence is a single cell capable of producing a single, dormant spore in response to an unfavourable environment, as seen for some bacteria, amoebzoa and yeasts. Here, near the upper extreme of environmental dependence, reproductive competence must be tightly integrated, if not synonymous, with the cell cycle: a phase of nutrient acquisition and biomass accumulation, followed by DNA replication and differentiation of a distinct cell type. Tracing this ontogenic phase back in time with evolutionary developmental studies would reveal whether competence in multicellular organisms shares genetic heritage with the cell cycle. In this regard, the finding that *ras* activity controls developmental transitions, nuclear division and polarity in *Aspergilli* [50,99,123], and the essentiality of the CSN in regulation of both competence and the cell cycle warrants further attention [123,124].

Beyond evolutionarily related reproductive competence states where conservation of underlying genetic architecture could be expected, other thresholds may share only conceptual homology due to emergent organization [125]. In other words, competence thresholds are simply necessary for efficient multicellular development, as seen most clearly in the multiple competence states delineated during development of *D. discoideum* from independent, single cells through a multicellular, motile aggregated stage to construction of a fruiting body for dispersal of dormant spores. Achievement of each state requires coordinated signalling, and competence in at least one case is as simple as expression, or not, of a single gene, the receptor for cAMP. So competence may be adaptive, in the sense that coordinated development is obviously adaptive, but competence systems in different organisms may be adapted to their current respective niches to very different degrees.

In the case of fungi, there is little research on competence generally, and certainly no evidence that timing is an

ecologically relevant, selected trait. This is in contrast to, for example, the widespread phenomenon of conidial autoinhibition (density-dependent inhibition of germination), which can at least be suggested as a plausible mechanism to promote dispersal and reduce competition [126], or circadian rhythmicity for conidiation [127].

If transition points between discrete developmental phases are optimized thresholds, whether of nutrient acquisition or environmental sensing (or hormones reporting these cues), questions of what these thresholds physically represent, what deviation from the optimum means for the organism, the population genetic or epigenetic variation for these traits and the relationship to organism ecology remain to be addressed. Answers to these questions in experimentally tractable fungi would require broad population genetic research looking for variation in competence timing among wild-isolates, mechanistic study of the effects of manipulating competence timing on reproductive output, and biochemical and genetic study to determine the factors of relevance for competence timing and the networks underlying their production.

5. Summary

Interplay between nutrient acquisition and parental nutrition, coordinated by chemical messengers, is a recurrent theme in development of plants and organisms spanning *Dictyostelids* to mammals. Discrete life phases in Unikonts may trace their roots to early spore-bearing species subject to a strict competence threshold integrated with the cell cycle. The existence of reproductive competence in fungi is well established, though not well appreciated, and its genetic and epigenetic basis, taxonomic distribution and current ecological relevance, if any, are poorly understood. Recent transcriptional profiling of *A. fumigatus* has revealed significant transcriptional rewiring associated with competence acquisition [128], and early biochemical work in *A. nidulans* shows pleiotropic change in membrane transport, intracellular iron availability and other systems. This clearly necessitates controlling for competence state in experimental studies. With amoebzoa such as the *Dictyostelids*, study of competence in a range of fungal species is required to establish its generality and relevance to other more elaborate eukaryotic systems.

Acknowledgements. We thank Hannah Robertson and Len Kelly for use of their critical faculties, David Axelrod, Yasuo Maeda, Bill Loomis and Richard Gomer for helpful discussions, and apologize to the many authors whose work we were unable to cite due to space limitations.

Funding statement. This work was supported by a grant from the Australian Research Council to A.A. and an Australian Postgraduate Award to L.M.N.

References

1. Waddington C. 1936 The origin of competence for lens formation in the Amphibia. *J. Exp. Biol.* **13**, 86.
2. Waddington CH. 1940 *Organisers and genes*. Cambridge, UK: Cambridge University Press.
3. Gross PR. 1968 Biochemistry of differentiation. *Annu. Rev. Biochem.* **37**, 631–660. (doi:10.1146/annurev.bi.37.070168.003215)
4. Slack JM. 2002 Conrad Hal Waddington: the last Renaissance biologist? *Nat. Rev. Genet.* **3**, 889–895. (doi:10.1038/nrg933)
5. Waddington C. 1932 Experiments on the development of chick and duck embryos, cultivated *in vitro*. *Phil. Trans. R. Soc. Lond. B* **221**, 179–230. (doi:10.1098/rstb.1932.0003)
6. Wolpert L. 1992 Gastrulation and the evolution of development. *Development* **1992**, 7–13.
7. Rudel D, Sommer RJ. 2003 The evolution of developmental mechanisms. *Dev. Biol.* **264**, 15–37. (doi:10.1016/S0012-1606(03)00353-1)
8. Davidson EH, McClay DR, Hood L. 2003 Regulatory gene networks and the properties of the

- developmental process. *Proc. Natl Acad. Sci. USA* **100**, 1475–1480. (doi:10.1073/pnas.0437746100)
9. Gluckman PD, Hanson MA. 2006 Evolution, development and timing of puberty. *Trends Endocrinol. Metab.* **17**, 7–12. (doi:10.1016/j.tem.2005.11.006)
 10. Bergonzi S, Albani MC. 2011 Reproductive competence from an annual and a perennial perspective. *J. Exp. Bot.* **62**, 4415–4422. (doi:10.1093/jxb/err192)
 11. Poethig RS. 2010 The past, present, and future of vegetative phase change. *Plant Physiol.* **154**, 541–544. (doi:10.1104/pp.110.161620)
 12. Axelrod DE. 1972 Kinetics of differentiation of conidiophores and conidia by colonies of *Aspergillus nidulans*. *J. Gen. Microbiol.* **73**, 181–184. (doi:10.1099/00221287-73-1-181)
 13. Casselton L, Zolan M. 2002 The art and design of genetic screens: filamentous fungi. *Nat. Rev. Genet.* **3**, 683–697. (doi:10.1038/nrg889)
 14. Axelrod DE, Gealt M, Pastushok M. 1973 Gene control of developmental competence in *Aspergillus nidulans*. *Dev. Biol.* **34**, 9–15. (doi:10.1016/0012-1606(73)90335-7)
 15. Trinci AP. 1969 A kinetic study of the growth of *Aspergillus nidulans* and other fungi. *J. Gen. Microbiol.* **57**, 11–24. (doi:10.1099/00221287-57-1-11)
 16. Lafon A, Seo JA, Han KH, Yu JH, d'Enfert C. 2005 The heterotrimeric G-protein GanB(α)-SfaD(β)-GpgA(γ) is a carbon source sensor involved in early cAMP-dependent germination in *Aspergillus nidulans*. *Genetics* **171**, 71–80. (doi:10.1534/genetics.105.040584)
 17. Fillinger S, Chaveroche MK, Shimizu K, Keller N, d'Enfert C. 2002 cAMP and ras signalling independently control spore germination in the filamentous fungus *Aspergillus nidulans*. *Mol. Microbiol.* **44**, 1001–1016. (doi:10.1046/j.1365-2958.2002.02933.x)
 18. Oshero N, May G. 2000 Conidial germination in *Aspergillus nidulans* requires RAS signaling and protein synthesis. *Genetics* **155**, 647–656.
 19. Fiddy C, Trinci AP. 1976 Mitosis, septation, branching and the duplication cycle in *Aspergillus nidulans*. *J. Gen. Microbiol.* **97**, 169–184. (doi:10.1099/00221287-97-2-169)
 20. Momany M, Taylor I. 2000 Landmarks in the early duplication cycles of *Aspergillus fumigatus* and *Aspergillus nidulans*: polarity, germ tube emergence and septation. *Microbiology* **146**, 3279–3284.
 21. Ugalde U. 2006 Autoregulatory signals in mycelial fungi. In *The Mycota: growth, differentiation and sexuality* (eds U Kües, R Fischer), pp. 203–213. Berlin, Germany: Springer.
 22. Meyer V, Arentshorst M, Flitter SJ, Nitsche BM, Kwon MJ, Reynaga-Pena CG, Bartnicki-Garcia S, van den Hondel CA, Ram AF. 2009 Reconstruction of signaling networks regulating fungal morphogenesis by transcriptomics. *Eukaryot. Cell* **8**, 1677–1691. (doi:10.1128/EC.00050-09)
 23. Penalva MA, Tilburn J, Bignell E, Arst Jr HN. 2008 Ambient pH gene regulation in fungi: making connections. *Trends Microbiol.* **16**, 291–300. (doi:10.1016/j.tim.2008.03.006)
 24. Lengeler KB, Davidson RC, D'Souza C, Harashima T, Shen WC, Wang P, Pan X, Waugh M, Heitman J. 2000 Signal transduction cascades regulating fungal development and virulence. *Microbiol. Mol. Biol. Rev.* **64**, 746–785. (doi:10.1128/MMBR.64.4.746-785.2000)
 25. Aguirre J. 1993 Spatial and temporal controls of the *Aspergillus* brlA developmental regulatory gene. *Mol. Microbiol.* **8**, 211–218. (doi:10.1111/j.1365-2958.1993.tb01565.x)
 26. Han S, Adams TH. 2001 Complex control of the developmental regulatory locus brlA in *Aspergillus nidulans*. *Mol. Genet. Genomics* **266**, 260–270. (doi:10.1007/s004380100552)
 27. Prade RA, Timberlake WE. 1993 The *Aspergillus nidulans* brlA regulatory locus consists of overlapping transcription units that are individually required for conidiophore development. *EMBO J.* **12**, 2439–2447.
 28. Kasuga T, Glass N. 2008 Dissecting colony development of *Neurospora crassa* using mRNA profiling and comparative genomics approaches. *Eukaryot. Cell* **7**, 1549–1564. (doi:10.1128/EC.00195-08)
 29. Greenwald CJ, Kasuga T, Glass NL, Shaw BD, Ebbole DJ, Wilkinson HH. 2010 Temporal and spatial regulation of gene expression during asexual development of *Neurospora crassa*. *Genetics* **186**, 1217–1230. (doi:10.1534/genetics.110.121780)
 30. Vinck A, Terlou M, Pestman WR, Martens EP, Ram AF, van den Hondel CA, Wosten HA. 2005 Hyphal differentiation in the exploring mycelium of *Aspergillus niger*. *Mol. Microbiol.* **58**, 693–699. (doi:10.1111/j.1365-2958.2005.04869.x)
 31. Masai K, Maruyama J, Sakamoto K, Nakajima H, Akito O. 2006 Square-plate culture method allows detection of differential gene expression and screening of novel, region-specific genes in *Aspergillus oryzae*. *Appl. Microbiol. Biotechnol.* **71**, 881–891. (doi:10.1007/s00253-006-0429-z)
 32. Murray SL, Hynes MJ. 2010 Metabolic and developmental effects resulting from deletion of the citA gene encoding citrate synthase in *Aspergillus nidulans*. *Eukaryot. Cell* **9**, 656–666. (doi:10.1128/EC.00373-09)
 33. Marhouf JF, Adams TH. 1995 Identification of developmental regulatory genes in *Aspergillus nidulans* by overexpression. *Genetics* **139**, 537–547.
 34. Adams TH, Timberlake WE. 1990 Developmental repression of growth and gene expression in *Aspergillus*. *Proc. Natl Acad. Sci. USA* **87**, 5405–5409. (doi:10.1073/pnas.87.14.5405)
 35. Gealt MA, Axelrod DE. 1974 Coordinate regulation of enzyme inducibility and developmental competence in *Aspergillus nidulans*. *Dev. Biol.* **41**, 224–232. (doi:10.1016/0012-1606(74)90301-7)
 36. Clutterbuck AJ. 1975 Cyclic AMP levels during growth and conidiation. *Aspergillus Newsl.* **12**, 13–15.
 37. d'Enfert C, Bonini BM, Zapella PD, Fontaine T, Chaveroche MK, da Silva AM, Terenzi HF. 1999 cAMP-dependent neutral trehalases catalyze intracellular trehalose breakdown during spore germination in *Aspergillus nidulans* and *Neurospora crassa*. *Fungal Genet. Newsl.* **465**, 139.
 38. Lee N, D'Souza CA, Kronstad JW. 2003 Of smuts, blasts, mildews, and blights: cAMP signaling in phytopathogenic fungi. *Annu. Rev. Phytopathol.* **41**, 399–427. (doi:10.1146/annurev.phyto.41.052002.095728)
 39. Li G, Zhou X, Xu JR. 2012 Genetic control of infection-related development in *Magnaporthe oryzae*. *Curr. Opin. Microbiol.* **15**, 678–684. (doi:10.1016/j.mib.2012.09.004)
 40. Lafon A, Han KH, Seo JA, Yu JH, d'Enfert C. 2006 G-protein and cAMP-mediated signaling in *Aspergilli*: a genomic perspective. *Fungal Genet. Biol.* **43**, 490–502. (doi:10.1016/j.fgb.2006.02.001)
 41. Pastushok M, Axelrod DE. 1976 Effect of glucose, ammonium and media maintenance on the time of conidiophore initiation by surface colonies of *Aspergillus nidulans*. *J. Gen. Microbiol.* **94**, 221–224. (doi:10.1099/00221287-94-1-221)
 42. Hall NE, Axelrod DE. 1978 Sporulation competence in *Aspergillus nidulans*: a role for iron in development. *Cell Differ.* **7**, 73–82. (doi:10.1016/0045-6039(78)90008-8)
 43. Kurtz MB, Champe SP. 1979 Genetic control of transport loss during development of *Aspergillus nidulans*. *Dev. Biol.* **70**, 82–88. (doi:10.1016/0012-1606(79)90008-3)
 44. Kurtz M. 1980 Regulation of fructose transport during growth of *Aspergillus nidulans*. *J. Gen. Microbiol.* **118**, 389–396. (doi:10.1099/00221287-118-2-389)
 45. Wei N, Deng XW. 1992 COP9: a new genetic locus involved in light-regulated development and gene expression in *Arabidopsis*. *Plant Cell* **4**, 1507–1518. (doi:10.1105/tpc.4.12.1507)
 46. von Arnim AG, Osterlund MT, Kwok SF, Deng XW. 1997 Genetic and developmental control of nuclear accumulation of COP1, a repressor of photomorphogenesis in *Arabidopsis*. *Plant Physiol.* **114**, 779–788. (doi:10.1104/pp.114.3.779)
 47. Butnick NZ, Yager LN, Kurtz MB, Champe SP. 1984 Genetic analysis of mutants of *Aspergillus nidulans* blocked at an early stage of sporulation. *J. Bacteriol.* **160**, 541–545.
 48. Yager LN, Kurtz MB, Champe SP. 1982 Temperature-shift analysis of conidial development in *Aspergillus nidulans*. *Dev. Biol.* **93**, 92–103. (doi:10.1016/0012-1606(82)90242-1)
 49. Lewis C, Champe SP. 1995 A pre-induction sporulation gene from *Aspergillus nidulans*. *Microbiology* **141**, 1821–1828. (doi:10.1099/13500872-141-8-1821)
 50. Som T, Kolaparthi VS. 1994 Developmental decisions in *Aspergillus nidulans* are modulated by ras activity. *Mol. Cell. Biol.* **14**, 5333–5348.
 51. Steyaert JM, Weld RJ, Stewart A. 2010 Ambient pH intrinsically influences *Trichoderma* conidiation and colony morphology. *Fungal Biol.* **114**, 198–208. (doi:10.1016/j.funbio.2009.12.004)

52. Gressel J, Galun E. 1967 Morphogenesis in *Trichoderma*: photoinduction and RNA. *Dev. Biol.* **15**, 575–598. (doi:10.1016/0012-1606(67)90054-1)
53. Hadley G, Harrold C. 1958 The sporulation of *Penicillium notatum* Westling in submerged liquid culture. II. The initial sporulation phase. *J. Exp. Bot.* **9**, 418–425. (doi:10.1093/jxb/9.3.418)
54. Keller NP, Turner G, Bennett JW. 2005 Fungal secondary metabolism: from biochemistry to genomics. *Nat. Rev. Microbiol.* **3**, 937–947. (doi:10.1038/nrmicro1286)
55. Roncal T, Ugalde U. 2003 Conidiation induction in *Penicillium*. *Res. Microbiol.* **154**, 539–546. (doi:10.1016/S0923-2508(03)00168-2)
56. Foster J, McDaniel L, Woodruff H, Stokes J. 1945 Microbiological aspects of penicillin: V. conidiospore formation in submerged cultures of *Penicillium notatum*. *J. Bacteriol.* **50**, 365.
57. Hadley G, Harrold C. 1958 The sporulation of *Penicillium notatum* Westling in submerged liquid culture. I. The effect of calcium and nutrients on sporulation intensity. *J. Exp. Bot.* **9**, 408–417. (doi:10.1093/jxb/9.3.408)
58. Morton AG, England DJF, Towler DA. 1958 The physiology of sporulation in *Penicillium griseofulvum* Dierckx. *Trans. Br. Mycol. Soc.* **41**, 39–51. (doi:10.1016/S0007-1536(58)80004-2)
59. Roncal T, Cordobes S, Sterner O, Ugalde U. 2002 Conidiation in *Penicillium cyclopium* is induced by conidiogenone, an endogenous diterpene. *Eukaryot. Cell* **1**, 823–829. (doi:10.1128/EC.1.5.823-829.2002)
60. Lawson EJ, Poethig RS. 1995 Shoot development in plants: time for a change. *Trends Genet.* **11**, 263–268. (doi:10.1016/S0168-9525(00)89072-1)
61. Kerstetter RA, Poethig RS. 1998 The specification of leaf identity during shoot development. *Annu. Rev. Cell Dev. Biol.* **14**, 373–398. (doi:10.1146/annurev.cellbio.14.1.373)
62. Poethig RS. 2003 Phase change and the regulation of developmental timing in plants. *Science* **301**, 334–336. (doi:10.1126/science.1085328)
63. Ahmed ML, Ong KK, Dunger DB. 2009 Childhood obesity and the timing of puberty. *Trends Endocrinol. Metab.* **20**, 237–242. (doi:10.1016/j.tem.2009.02.004)
64. DiVall SA, Radovick S. 2008 Pubertal development and menarche. *Ann. NY Acad. Sci.* **1135**, 19–28. (doi:10.1196/annals.1429.026)
65. Roa J, Garcia-Galiano D, Castellano JM, Gaytan F, Pinilla L, Tena-Sempere M. 2010 Metabolic control of puberty onset: new players, new mechanisms. *Mol. Cell Endocrinol.* **324**, 87–94. (doi:10.1016/j.mce.2009.12.018)
66. Vandenberg JG, Drickamer LC, Colby DR. 1972 Social and dietary factors in the sexual maturation of female mice. *J. Reprod. Fertil.* **28**, 397–405. (doi:10.1530/jrf.0.0280397)
67. Short RV. 1976 The evolution of human reproduction. *Proc. R. Soc. Lond. B* **195**, 3–24. (doi:10.1098/rspb.1976.0095)
68. Brink RA. 1962 Phase change in higher plants and somatic cell heredity. *Q. Rev. Biol.* **37**, 1–22. (doi:10.1086/403567)
69. Mauseth J. 2013 Cephalia: juvenile/adult phase change in Cacti. See <http://www.sbs.utexas.edu/mauseth/>.
70. Proveniers M. 2013 Sugars speed up the circle of life. *Elife* **2**, e00625. (doi:10.7554/eLife.00625)
71. Baurle I, Dean C. 2006 The timing of developmental transitions in plants. *Cell* **125**, 655–664. (doi:10.1016/j.cell.2006.05.005)
72. Evans MM, Poethig RS. 1995 Gibberellins promote vegetative phase change and reproductive maturity in maize. *Plant Physiol.* **108**, 475–487. (doi:10.1104/pp.108.2.475)
73. Telfer A, Bollman KM, Poethig RS. 1997 Phase change and the regulation of trichome distribution in *Arabidopsis thaliana*. *Development* **124**, 645–654.
74. Wang JW, Park MY, Wang LJ, Koo Y, Chen XY, Weigel D, Poethig RS. 2011 miRNA control of vegetative phase change in trees. *PLoS Genet.* **7**, e1002012. (doi:10.1371/journal.pgen.1002012)
75. Earley K, Smith M, Weber R, Gregory B, Poethig R. 2010 An endogenous F-box protein regulates ARGONAUTE1 in *Arabidopsis thaliana*. *Silence* **1**, 15. (doi:10.1186/1758-907X-1-15)
76. Poethig RS. 2009 Small RNAs and developmental timing in plants. *Curr. Opin. Genet. Dev.* **19**, 374–378. (doi:10.1016/j.gde.2009.06.001)
77. Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS. 2009 The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* **138**, 750–759. (doi:10.1016/j.cell.2009.06.031)
78. Wang JW, Czech B, Weigel D. 2009 miR156-regulated SPL transcription factors define an endogenous flowering pathway in *Arabidopsis thaliana*. *Cell* **138**, 738–749. (doi:10.1016/j.cell.2009.06.014)
79. Yang L, Xu M, Koo Y, He J, Poethig RS. 2013 Sugar promotes vegetative phase change in *Arabidopsis thaliana* by repressing the expression of MIR156A and MIR156C. *Elife* **2**, e00260. (doi:10.7554/eLife.00260)
80. Yu S *et al.* 2012 Gibberellin regulates the *Arabidopsis* floral transition through miR156-targeted SQUAMOSA promoter binding-like transcription factors. *Plant Cell* **24**, 3320–3332. (doi:10.1105/tpc.112.101014)
81. McLaughlin DJ, Hibbett DS, Lutzoni F, Spatafora JW, Vilgalys R. 2009 The search for the fungal tree of life. *Trends Microbiol.* **17**, 488–497. (doi:10.1016/j.tim.2009.08.001)
82. Koonin EV. 2010 The origin and early evolution of eukaryotes in the light of phylogenomics. *Genome Biol.* **11**, 209. (doi:10.1186/gb-2010-11-5-209)
83. Schaap P *et al.* 2006 Molecular phylogeny and evolution of morphology in the social amoebas. *Science* **314**, 661–663. (doi:10.1126/science.1130670)
84. Kaiser D. 2003 Coupling cell movement to multicellular development in myxobacteria. *Nat. Rev. Microbiol.* **1**, 45–54. (doi:10.1038/nrmicro733)
85. Rosenberg E, DeLong E, Lory S, Stackebrandt E, Thompson F. 2013 *The prokaryotes: prokaryotic communities and ecophysiology*. Berlin, Germany: Springer.
86. Harris BZ, Kaiser D, Singer M. 1998 The guanosine nucleotide (p)ppGpp initiates development and A-factor production in *Myxococcus xanthus*. *Genes Dev.* **12**, 1022–1035. (doi:10.1101/gad.12.7.1022)
87. Bakthavatsalam D, Brazill D, Gomer RH, Eichinger L, Rivero F, Noegel AA. 2007 A G protein-coupled receptor with a lipid kinase domain is involved in cell-density sensing. *Curr. Biol.* **17**, 892–897. (doi:10.1016/j.cub.2007.04.029)
88. Gao T *et al.* 2007 A cell number-counting factor regulates levels of a novel protein, SslA, as part of a group size regulation mechanism in *Dictyostelium*. *Eukaryot. Cell* **6**, 1538–1551. (doi:10.1128/EC.00169-07)
89. Katoh M, Chen G, Roberge E, Shaulsky G, Kuspa A. 2007 Developmental commitment in *Dictyostelium discoideum*. *Eukaryot. Cell* **6**, 2038. (doi:10.1128/EC.00223-07)
90. Oohata AA, Fukuzawa M, Hotta R, Nakagawa M, Niwa M, Takaya Y. 2009 Differentiation inducing factors in *Dictyostelium discoideum*: a novel low molecular factor functions at an early stage(s) of differentiation. *Dev. Growth Differ.* **51**, 743–752. (doi:10.1111/j.1440-169X.2009.01133.x)
91. Okamoto K. 1985 Acquisition process of differentiation competence in *Dictyostelium discoideum*. *Dev. Biol.* **107**, 198–205. (doi:10.1016/0012-1606(85)90387-2)
92. Maeda Y. 2011 Cell-cycle checkpoint for transition from cell division to differentiation. *Dev. Growth Differ.* **53**, 463–481. (doi:10.1111/j.1440-169X.2011.01264.x)
93. Loomis WF, Shaulsky G. 2011 Developmental changes in transcriptional profiles. *Dev. Growth Differ.* **53**, 567–575. (doi:10.1111/j.1440-169X.2010.01241.x)
94. Bonner JT, Joyner BD, Moore AA, Suthers HB, Swanson JA. 1985 Successive asexual life cycles of starved amoebae in the cellular slime mould, *Dictyostelium mucoroides* var. *stoloniferum*. *J. Cell. Sci.* **77**, 19–26.
95. Bull AT, Trinci AP. 1977 The physiology and metabolic control of fungal growth. *Adv. Microb. Physiol.* **15**, 1–84. (doi:10.1016/S0065-2911(08)60314-8)
96. Rodriguez-Urra AB, Jimenez C, Duenas M, Ugalde U. 2009 Bicarbonate gradients modulate growth and colony morphology in *Aspergillus nidulans*. *FEMS Microbiol. Lett.* **300**, 216–221. (doi:10.1111/j.1574-6968.2009.01780.x)
97. Semighini CP, Harris SD. 2008 Regulation of apical dominance in *Aspergillus nidulans* hyphae by reactive oxygen species. *Genetics* **179**, 1919–1932. (doi:10.1534/genetics.108.089318)
98. Rittenour W, Si H, Harris S. 2009 Hyphal morphogenesis in *Aspergillus nidulans*. *Fungal Biol. Rev.* **23**, 20–29. (doi:10.1007/s10267-009-08.001)
99. Harispe L, Portela C, Scacciochio C, Penalva MA, Gorfinkiel L. 2008 Ras GTPase-activating protein regulation of actin cytoskeleton and hyphal polarity

- in *Aspergillus nidulans*. *Eukaryot. Cell* **7**, 141–153. (doi:10.1128/EC.00346-07)
100. Araujo-Bazán L, Peñalva MA, Espeso EA. 2008 Preferential localization of the endocytic internalization machinery to hyphal tips underlies polarization of the actin cytoskeleton in *Aspergillus nidulans*. *Mol. Microbiol.* **67**, 891–905. (doi:10.1111/j.1365-2958.2007.06102.x)
101. Roncal T, Ugalde UO, Irastorza A. 1993 Calcium-induced conidiation in *Penicillium cyclopium*: calcium triggers cytosolic alkalization at the hyphal tip. *J. Bacteriol.* **175**, 879–886.
102. Robson G, Prebble E, Rickers A, Hosking S, Denning D, Trinci A, Robertson W. 1996 Polarized growth of fungal hyphae is defined by an alkaline pH gradient. *Fungal Genet. Biol.* **20**, 289–298. (doi:10.1006/fghi.1996.0043)
103. Hense BA, Kuttler C, Muller J, Rothballer M, Hartmann A, Kreft JU. 2007 Does efficiency sensing unify diffusion and quorum sensing? *Nat. Rev. Microbiol.* **5**, 230–239. (doi:10.1038/nrmicro1600)
104. Brodhun F, Feussner I. 2011 Oxylipins in fungi. *FEBS J.* **278**, 1047–1063. (doi:10.1111/j.1742-4658.2011.08027.x)
105. Herrero-García E, Garzia A, Cordobes S, Espeso EA, Ugalde U. 2011 8-Carbon oxylipins inhibit germination and growth, and stimulate aerial conidiation in *Aspergillus nidulans*. *Fungal Biol.* **115**, 393–400. (doi:10.1016/j.funbio.2011.02.005)
106. Lee DS, Nioche P, Hamberg M, Raman CS. 2008 Structural insights into the evolutionary paths of oxylipin biosynthetic enzymes. *Nature* **455**, 363–368. (doi:10.1038/nature07307)
107. Tsitsigiannis DI, Keller NP. 2007 Oxylipins as developmental and host-fungal communication signals. *Trends Microbiol.* **15**, 109–118. (doi:10.1016/j.tim.2007.01.005)
108. Rodríguez-Urra AB, Jiménez C, Nieto MI, Rodríguez J, Hayashi H, Ugalde U. 2012 Signaling the induction of sporulation involves the interaction of two secondary metabolites in *Aspergillus nidulans*. *ACS Chem. Biol.* **7**, 599–606. (doi:10.1021/cb200455u)
109. Sandhu J, Ben-Shlomo Y, Cole TJ, Holly J, Davey Smith G. 2006 The impact of childhood body mass index on timing of puberty, adult stature and obesity: a follow-up study based on adolescent anthropometry recorded at Christ's Hospital (1936–1964). *Int. J. Obes.* **30**, 14–22. (doi:10.1038/sj.ijo.0803156)
110. Lovett Doust J, Lovett Doust L. 1988 *Plant reproductive ecology: patterns and strategies*. New York, NY: Oxford University Press.
111. Plaistow SJ, Lapsley CT, Beckerman AP, Benton TG. 2004 Age and size at maturity: sex, environmental variability and developmental thresholds. *Proc. R. Soc. Lond. B* **271**, 919–924. (doi:10.1098/rsbp.2004.2682)
112. Callier V, Nijhout HF. 2011 Control of body size by oxygen supply reveals size-dependent and size-independent mechanisms of molting and metamorphosis. *Proc. Natl Acad. Sci. USA* **108**, 14 664–14 669. (doi:10.1073/pnas.1106556108)
113. Davidowitz G, D'Amico LJ, Nijhout HF. 2003 Critical weight in the development of insect body size. *Evol. Dev.* **5**, 188–197. (doi:10.1046/j.1525-142X.2003.03026.x)
114. Beadle G, Tatum E, Clancy C. 1938 Food level in relation to rate of development and eye pigmentation in *Drosophila melanogaster*. *Biol. Bull.* **75**, 447–462. (doi:10.2307/1537573)
115. Whalen MC. 2005 Host defence in a developmental context. *Mol. Plant Pathol.* **6**, 347–360. (doi:10.1111/j.1364-3703.2005.00286.x)
116. Kearsley MJC, Whitham TG. 1989 Developmental changes in resistance to herbivory: implications for individuals and populations. *Ecology* **70**, 422–434. (doi:10.2307/1937547)
117. Hadfield MG, Carpizo-Etuarte EJ, del Carmien K, Nedved BT. 2001 Metamorphic competence, a major adaptive convergence in marine invertebrate larvae. *Am. Zool.* **41**, 1123–1131. (doi:10.1668/0003-1569(2001)041[1123:MCAMAC]2.0.CO;2)
118. Nielsen C. 1998 Origin and evolution of animal life cycles. *Biol. Rev.* **73**, 125–155. (doi:10.1017/S0006323197005136)
119. Hodin J. 2006 Expanding networks: signaling components in and a hypothesis for the evolution of metamorphosis. *Integr. Comp. Biol.* **46**, 719–742. (doi:10.1093/icb/icl038)
120. Tennessen J, Thummel C. 2011 Coordinating growth and maturation: insights from *Drosophila*. *Curr. Biol.* **21**, R750–R757. (doi:10.1016/j.cub.2011.06.033)
121. Hölldobler B, Wilson E. 1990 *The ants*. Cambridge, MA: Harvard University Press.
122. Fortwendel JR, Panepinto JC, Seitz AE, Askew DS, Rhodes JC. 2004 *Aspergillus fumigatus* rasA and rasB regulate the timing and morphology of asexual development. *Fungal Genet. Biol.* **41**, 129–139. (doi:10.1016/j.fgb.2003.10.004)
123. Wei N, Serino G, Deng XW. 2008 The COP9 signalosome: more than a protease. *Trends Biochem. Sci.* **33**, 592–600. (doi:10.1016/j.tibs.2008.09.004)
124. Adams TH, Hide WA, Yager LN, Lee BN. 1992 Isolation of a gene required for programmed initiation of development by *Aspergillus nidulans*. *Mol. Cell. Biol.* **12**, 3827–3833.
125. Kauffman SA. 1993 *The origins of order: self-organization and selection in evolution*. New York, NY: Oxford University Press.
126. Leeder AC, Palma-Guerrero J, Glass NL. 2011 The social network: deciphering fungal language. *Nat. Rev. Microbiol.* **9**, 440–451. (doi:10.1038/nrmicro2580)
127. Dunlap JC, Loros JJ. 2006 How fungi keep time: circadian system in *Neurospora* and other fungi. *Curr. Opin. Microbiol.* **9**, 579–587. (doi:10.1016/j.mib.2006.10.008)
128. Sheppard DC, Doedt T, Chiang LY, Kim HS, Chen D, Nierman WC, Filler SG. 2005 The *Aspergillus fumigatus* StuA protein governs the up-regulation of a discrete transcriptional program during the acquisition of developmental competence. *Mol. Biol. Cell* **16**, 5866–5879. (doi:10.1091/mbc.E05-07-0617)